

LECTURE OUTLINE

1. Ways of reaching saturation

- Isobaric processes
 - Isobaric cooling (dew point temperature)
 - Isobaric and adiabatic cooling by evaporation of water (wet bulb temperature)
 - Adiabatic and isobaric mixing



WAYS OF REACHING SATURATION FORMATION AND DISSIPATION OF CLOUDS

For simplicity, we assume that clouds form in the atmosphere when the water vapor reaches the saturation value, and $f=100\%$ (in reality this value should be greater than 100%).

$$f = \frac{q_v}{q_s(p, T)}$$

An increase of relative humidity can be accomplished by:

- increasing the amount of water vapor in the air (i.e. increasing q_v); evaporation of water from a surface or via evaporation of rain falling through unsaturated air,
- cooling of the air (decrease of $q_s(p, T)$); isobaric cooling (e.g. radiative), adiabatic cooling of rising air,
- mixing of two unsaturated parcels of air.



THE COMBINED FIRST AND SECOND LAWS WILL BE APPLIED TO THE IDEALIZED THERMODYNAMIC REFERENCE PROCESSES ASSOCIATED WITH PHASE CHANGE OF WATER:

1. isobaric cooling
2. adiabatic isobaric processes
3. adiabatic and isobaric mixing




R&Y, Chapter 2

Salby, Chapter 5



A Short Course in Cloud Physics,
R.R. Rogers and M.K. Yau; R&Y

C&W, Chapter 6



Thermodynamics of Atmospheres
and Oceans,
J.A. Curry and P.J. Webster; C&W

ISOBARIC PROCESSES LEADING TO SATURATION OF THE AIR WITH WATER VAPOR

1. Isobaric cooling ($p=const, q_v=const$)
 - Dew point temperature
 - Isobaric cooling with condensation ($p=const, q_t=const$)
2. Isobaric and adiabatic cooling/heating by evaporation/condensation of water ($p=const, a\ source\ of\ water\ vapor\ / \ water$)
 - Wet bulb temperature
 - Equivalent temperature
3. Isobaric and adiabatic mixing ($p=const, q_t=const$)



ISOBARIC COOLING



UNIVERSITY
OF WARSAW

If vertical movement in the atmosphere is insignificant and a departure from the initial state is small, processes can be seen as **isobaric processes**.

If there is no condensation, the First Law of thermodynamics is:

$$\delta q = dh = c_p dT$$

c_p takes a value for the moist air $c_p = c_{pd}(1 + 0.87q_v)$, but can also be approximated by its dry air c_{pd} .

If the air is cooled, the relative humidity (f) will increase:

$$f = \frac{e}{e_s(T)} \cong \frac{q_v}{q_{vs}(p, T)}$$

- q_v does not change;
- temperature is decreasing therefore the saturated water vapor pressure (e_s) decreases
- because the process is isobaric ($p = const$) the same is true for q_s , i.e. q_s decreases.

Cooling leads to the saturation ($f=1$) and then condensation starts.

DEW POINT TEMPERATURE - T_d

The dew point T_d is defined as that temperature to which the system must be cooled **isobarically** to achieve saturation


The dew point temperature is unchanged during the isobaric process without condensation.



If saturation occurs below 0°C , that temperature is the **frost point T_f** .

The **dew point temperature** is defined as: $e = e_s(T_d)$ or $q_v = q_s(T_d)$, where e is the actual pressure of water vapor in the air at temperature T , $q_v = \text{const}$ is the actual mixing ratio.

T_d can be calculated from equation: $e = e_{s0} \exp \left[\frac{L_{lv}}{R_v} \left(\frac{1}{T_T} - \frac{1}{T_d} \right) \right]$



e is the actual pressure of water vapor in the air at temperature T

The dew point temperature unit is Kelvin.

The dew point temperature is not a measure of temperature, but a measure of humidity in the air (relative humidity, f).


We will integrate the Clausius-Clapeyron equation between temperatures T and T_d .

$$\frac{d \ln p}{dT} = \frac{L_{lv}}{R_v T^2}$$


$$\int_{e_s}^e d \ln p = \int_T^{T_d} \frac{L_{lv}}{R_v T^2} dT$$

$$\ln \left(\frac{e}{e_s} \right) = \ln f = \frac{L_{lv}}{R_v} \left(\frac{1}{T} - \frac{1}{T_d} \right) \quad \text{We assume } L_{lv} = \text{const.}$$

$$f = \exp \left[- \frac{L_{lv}}{R_v} \left(\frac{T - T_d}{T \cdot T_d} \right) \right]$$




$T - T_d$ is the dew point deficit or the dew point spread.



Dew point temperature: $e = e_{s0} \exp \left[\frac{L_{lv}}{R_v} \left(\frac{1}{T_0} - \frac{1}{T_d} \right) \right]$

The frost point temperature: $e = e_{s0} \exp \left[\frac{L_{iv}}{R_v} \left(\frac{1}{T_0} - \frac{1}{T_f} \right) \right]$

$T_f: e = e_{si}(T_f)$ or $q_v = q_{si}(T_f)$



ISOBARIC COOLING WITH CONDENSATION

If the air is cooled isobarically below the dew point temperature (T_d), the condensation starts.

First Law of thermodynamics : $dh = \delta q + vdp$

Enthalpy for two-component system: $dh = c_p dT + L_{lv} dq_v$

For isobaric process $dp=0$:
 $dh = \delta q$
 $\delta q = c_p dT + L_{lv} dq_v$

Let's assume that condensation occurs always at saturation state ($f=1$) :

- the specific humidity is equal to its value at saturation state ($q_v = q_s$),
- $q_t = q_s + q_l$

In a closed system q_t does not change, i.e. $dq_t = 0$, and $dq_l = -dq_s$.

The amount of water dq_l , that condenses during isobaric cooling:

$$dq_l = -dq_s \cong -\frac{\varepsilon}{p} de_s = -\frac{\varepsilon L_{lv} e_s}{p R_v T^2} dT = -\frac{\varepsilon^2 L_{lv} e_s}{p R T^2} dT$$

$$q \cong \varepsilon \frac{e_s}{p}$$

$$\frac{de_s}{dT} = \frac{L_{lv} e_s}{R_v T^2}$$

$$\varepsilon = \frac{R}{R_v}$$

In the First Law equation for an isobaric process

$$\delta q = c_p dT + L_{lv} dq_v$$

- we express dq_v as a function of dT (see previous slide):

$$dq_v = \frac{\varepsilon^2 L_{lv} e_s}{pRT^2} dT$$

to find how the temperature changes if heat δq is added:

$$\delta q = \left(c_p + \frac{\varepsilon^2 L_{lv}^2 e_s}{pRT^2} \right) dT$$

Before condensation occurs $\delta q = c_p dT$.

After condensation starts the temperature decreases slower in response to isobaric cooling. It is due to the release of latent heat of condensation.

$$dT = \frac{\delta q}{c_p + \frac{\varepsilon^2 L_{lv}^2 e_s}{pRT^2}}$$

In the First Law equation for an isobaric process

$$\delta q = c_p dT + L_{lv} dq_v$$

- we express dT as a function of dq_l (see previous slide):

$$dT = -\frac{pRT^2}{\varepsilon^2 L_{lv} e_s} dq_l$$

to find how much heat should be taken out of the system to condense a given amount of water dq_l .

$$\delta q = -c_p \frac{pRT^2}{\varepsilon^2 L_{lv} e_s} dq_l - L_{lv} dq_l$$

$$dq_l = -\left(\frac{\varepsilon^2 L_{lv} e_s}{c_p p R T^2 + \varepsilon^2 L_{lv}^2 e_s} \right) \delta q$$

The relation must be integrated numerically, because e_s depends on temperature.

Once condensation begins, the dew-point temperature decreases, since the water vapor mixing ratio is decreasing as the water is condensed.

Relative humidity remains constant, at $f = 1 \rightarrow T_d = T$ $\left(f = \exp \left[-\frac{L_{lv}}{R_v} \left(\frac{T - T_d}{T \cdot T_d} \right) \right] \right)$

Isobaric cooling is a primary formation mechanism for certain types of fog and stratus clouds.

The equations derived are equally applicable for isobaric heating. In this instance, an existing cloud or fog can be dissipated by evaporation that ensues from isobaric heating (e.g. solar radiation).

ADIABATIC AND ISOBARIC COOLING AND MOISTENING BY WATER EVAPORATION



UNIVERSITY
OF WARSAW

WET-BULB TEMPERATURE, T_w

Consider a system composed of unsaturated moist air plus rain falling through the air. Because the air is undersaturated, the rain will evaporate.

If there is no external heat sources ($\Delta q=0$), and the evaporation occurs isobarically ($dp=0$), the enthalpy is conserved ($dh=0$):

$$dh = c_p dT + L_{lv} dq_v$$

$$0 = c_{pd} dT - L_{lv} dq_l$$

$$dh = 0$$

$$dq_v = -dq_l$$

$c_p = c_{pd}(1 + 0.87q_v)$ or can be approximated as the dry-air value c_{pd}

If we allow just enough liquid water from the rain to evaporate so that the air becomes saturated ($q_v = q_s$), we can integrate the above equation:

- from the state where there is q_l of liquid water at temperature T
- to the state where all liquid evaporates ($q_l = 0$) and the temperature decreased to T_w .

$$c_p \int_T^{T_w} dT = \int_{q_l}^0 L_{lv} dq_l$$

q_l is the amount of water that must be evaporated to bring the air to saturation at temperature T_w .

During the evaporation process, latent heat is drawn from the atmosphere, and the final temperature, referred to as the **wet-bulb temperature, T_w** , is cooler than the original temperature.

The wet bulb temperature – temperature to which air may be cooled by evaporating water into it at constant pressure, until saturation is reached.

$$c_p \int_T^{T_w} dT = \int_{q_l}^0 L_{lv} dq_l$$

$$c_p(T_w - T) = -L_{lv}q_l, \quad q_l = q_s(T_w, p) - q_v$$

$$T_w = T - \frac{L_{lv}}{c_p} [q_s(T_w, p) - q_v]$$

Temperature dependence of $L_{lv}(T)$ has been neglected. Given q_v , T and p , this expression is implicit for T_w and must be solved numerically.


However, if T and T_w are given, then q_v is easily determined.

The wet bulb temperature T_w is measured with the wet bulb thermometer.

It consists of two thermometers one of which is surrounded by a wet cloth and is ventilated.



Assman's psychrometer




©The COMET Program

The wet-bulb temperature in the atmosphere is conservative with respect to evaporation of falling rain.

Calculations for given values of T and q_v show that $T_d < T_w < T$.

This can be shown graphically. Since e increases while T decreases during the approach to T_w , the Clausius-Clapeyron diagram looks like:



If ice is the evaporating phase, the ice-bulb temperature T_i can be analogously determined:

$$T_i = T - \frac{L_{iv}}{c_p} [q_s(T_i) - q_v]$$

It is easily shown that $T_i > T_w$

EQUIVALENT TEMPERATURE, T_e

Equivalent temperature (T_e) is defined as the temperature a sample of moist air would attain if all the moisture were condensed out at **constant pressure without any heat transfer** from/to the environment.

We use the same equation as before:

$$dh = c_p dT + L_{lv} dq_v$$

$$0 = c_p dT + L_{lv} dq_s$$

$$dh = 0$$

$$dq_v = dq_s$$

The equation is integrated

- from a state in which there is q_s of water vapor (saturated) at temperature T
- to a state in which all water vapor condensed ($q_s = 0$) and the temperature increased to T_e :

$$c_p \int_T^{T_e} dT = - \int_{q_s}^0 L_{lv} dq_s$$

$$c_p (T_e - T) = L_{lv} q_s$$

$$T_e = T + \frac{L_{lv} q_s}{c_p}$$

EQUIVALENT POTENTIAL TEMPERATURE

We defined the equivalent potential temperature that is conserved in wet adiabatic/pseudo-adiabatic processes.

$$\theta_e = T \left(\frac{p_0}{p} \right)^{\frac{R_e}{c_{pe}}} \Omega_e \exp \left(\frac{q_v L_{vl}}{c_{pe} T} \right)$$

$$\Omega_e = \left(\frac{R}{R_e} \right)^{\frac{R_e}{c_{pe}}} \left(\frac{e}{e_s} \right)^{-\frac{q_v R_v}{c_{pe}}}$$

$$c_{pe} = c_{pd} + q_t (c_l - c_{pd})$$

$$R_e = (1 - q_t) R_d$$

e/e_s defines the relative humidity. The term Ω_e is close to 0 and depends only very weakly on the thermodynamic state.

The expression for θ_e in the full form is complicated, but it is rarely used in this form for practical applications.

For many purposes far simpler expressions capture much of the essential physics.

$$\theta_e = \theta \exp \left(\frac{q_s L_{vl}}{c_p T} \right)$$

Equivalent temperature: $T_e = T + \frac{L_{vl}q_s}{c_p}$

Equivalent potential temperature $\theta_e \cong \theta \exp\left(\frac{q_s L_{vl}}{c_p T}\right)$

For small $L_{lv}q_s/c_p T$:

$$\exp\left(\frac{L_{vl}q_s}{c_p T}\right) \cong 1 + \frac{L_{vl}q_s}{c_p T} = \frac{1}{T} \left(T + \frac{L_{vl}q_s}{c_p}\right)$$

$$\theta_e = \frac{\theta}{T} \left(T + \frac{L_{lv}q_s}{c_p}\right) = \left(T + \frac{L_{lv}q_s}{c_p}\right) \cdot \left(\frac{p_0}{p}\right)^{R_d/c_{pd}} = T_e \left(\frac{p_0}{p}\right)^{R_d/c_{pd}}$$

$$\theta_e \cong T_e \left(\frac{p_0}{p}\right)^{R_d/c_{pd}}$$

It is a potential equivalent temperature.

ADIABATIC AND ISOBARIC MIXING



UNIVERSITY
OF WARSAW

In some conditions isobaric mixing of different undersaturated air masses can lead to fog formation (saturated mass).

Let's consider the isobaric mixing of two undersaturated air masses (Y_1 and Y_2) having different temperatures and water vapor content.

Assume that there is no condensation.



To describe the **adiabatic** and **isobaric** mixing we will use the First Law of thermodynamic (in enthalpy form).

$$dH = \delta q + Vdp$$

$$0 = dH \approx m_1 c_{pd} dT_1 + m_2 c_{pd} dT_2$$

dT_1 and dT_2 correspond to the temperature change that occurs in response of the mixing process.


We neglect the input of water vapor to the heat capacity.



We will integrate the equation from the initial state defined by temperatures T_1 and T_2 to the final state having the temperature T .

$$m_1 c_{pd}(T - T_1) + m_2 c_{pd}(T - T_2) \approx 0$$

$$T \approx \frac{m_1}{m_1 + m_2} T_1 + \frac{m_2}{m_1 + m_2} T_2$$



The total mass $m = m_1 + m_2$ of the system remains unchanged during the mixing process. The specific humidity of the mixture, q_v , is a weighted average of initial specific humidities q_{v1} and q_{v2} .

$$q_v = \frac{m_1}{m_1 + m_2} q_{v1} + \frac{m_2}{m_1 + m_2} q_{v2}$$


Partial pressures of water vapor mix also linearly because $q_v \approx \varepsilon e/p$, and the process is isobaric.

The values (e, T) at point Y' can be found by solving simultaneously the former equation and the Clausius Clapeyron equation.

$Y \rightarrow Y'$ proces is adiabatic and isobaric.

The amount of water condensed in this process is:

$$\Delta q_l = \frac{\varepsilon}{p} [e(Y) - e(Y')]$$



An example of cloud formation resulting from adiabatic and isobaric mixing:

- contrails
- mist/haze formed in exhaled air




Fig. 6.4 C&W

A plane flying at 200 mb ejects water vapor into the atmosphere at the temperature and water vapor pressure represented by point A (600K, 4mb). For atmospheric temperatures less than -47°C (226 K), the water vapor will condense forming condensation trails.